

## Systolic Hypertension in the Elderly Program (SHEP) Publications

Applegate WB, Pressel S, Wittes J, Luhr J, Shekelle RB, Camel GH, Greenlick MR, Hadley E, Moye L, Perry HM Jr., et al. Impact of the treatment of isolated systolic hypertension on behavioral variables. Results from the systolic hypertension in the elderly program. *Arch Intern Med* 1994;154:2154-2160.

Applegate WB, Davis BR, Black HR, Smith WM, Miller ST, Burlando AJ. Prevalence of postural hypotension at baseline in the Systolic Hypertension in the Elderly Program (SHEP) cohort. *J Am Geriatr Soc* 1991;39:1057-1064.

Bearden D, Allman R, McDonald R, Miller S, Pressel S, Petrovitch H. Age, race, and gender variation in the utilization of coronary artery bypass surgery and angioplasty in SHEP. SHEP Cooperative Research Group. Systolic Hypertension in the Elderly Program. *J Am Geriatr Soc* 1994;42:1143-1149.

Black HR, Unger D, Burlando A, Wright JC, Pressel SL, Allen R, McDonald RH, Surath H. Systolic Hypertension in the Elderly Program (SHEP). Part 6: Baseline physical examination findings. *Hypertension* 1991;17(Suppl 2):77-101.

Borhani NO, Applegate WB, Cutler JA, Davis BR, Furberg CD, Lakatos E, Page L, Perry HM, Smith WM, Probstfield JL. Systolic Hypertension in the Elderly Program (SHEP). Part 1: Rationale and design. *Hypertension* 1991;17(Suppl 2):2-15.

Brittain E, Palensky J, Blood J, Wittes J. Blinded subjective rankings as a method of assessing treatment effect: a large sample example from the Systolic Hypertension in the Elderly Program (SHEP). *Stat Med* 1997;16:681-693.

Curb JD, Pressel SL, Cutler JA, Savage PJ, Applegate WB, Black H, Camel G, Davis BR, Frost PH, Gonzalez N, Guthrie G, Oberman A, Rutan GH, Stamler J. Effect of diuretic-based antihypertensive treatment on cardiovascular disease risk in older diabetic patients with isolated systolic hypertension. Systolic Hypertension in the Elderly Program Cooperative Research Group [Published erratum appears in *JAMA* 1997;277:1356]. *JAMA* 1996;276:1886-1892.

Davis BR, Vogt T, Frost PH, Burlando A, Cohen J, Wilson A, Brass LM, Frishman W, Price T, Stamler J. Risk factors for stroke and type of stroke in persons with isolated systolic hypertension. Systolic Hypertension in the Elderly Program Cooperative Research Group. *Stroke* 1998;29:1333-1340.

Davis BR, Wittes J, Pressel S, Berge KG, Hawkins CM, Lakatos E, Moye LA, Probstfield JL. Statistical considerations in monitoring the Systolic Hypertension in the Elderly Program (SHEP). *Control Clin Trials* 1993;14:350-361.

Di Bari M, Pahor M, Franse LV, Shorr RI, Wan JY, Ferrucci L, Somes GW, Applegate WB. Dementia and disability outcomes in large hypertension trials: lessons learned from the

Systolic Hypertension in the Elderly Program (SHEP) trial. *Am J Epidemiol* 2001;153:72-78.

Franse LV, Pahor M, Di Bari M, Somes GW, Cushman WC, Applegate WB. Hypokalemia associated with diuretic use and cardiovascular events in the Systolic Hypertension in the Elderly Program. *Hypertension* 2000;35:1025-1030.

Franse LV, Pahor M, Di Bari M, Shorr RI, Wan JY, Somes GW, Applegate WB. Serum uric acid, diuretic treatment and risk of cardiovascular events in the Systolic Hypertension in the Elderly Program (SHEP). *J Hypertens* 2000;18:1149-1154.

Frost PH, Davis BR, Burlando AJ, Curb JD, Guthrie GP Jr., Isaacsohn JL, Wassertheil-Smoller S, Wilson AC, Stamler J. Serum lipids and incidence of coronary heart disease. Findings from the Systolic Hypertension in the Elderly Program (SHEP). *Circulation* 1996;94:2381-2388.

Frost PH, Davis BR, Burlando AJ, Curb JD, Guthrie GP, Isaacsohn JL, Wassertheil-Smoller S, Wilson AC, Stamler J. Coronary heart disease risk factors in men and women aged 60 years and older: findings from the Systolic Hypertension in the Elderly Program. *Circulation* 1996;94:26-34.

Hall WD, Davis BR, Frost P, Hoffmeier M, O'Brien JE, Pace S, Page L, Schneider KA, Stamler J. Systolic Hypertension in the Elderly Program (SHEP). Part 7: Baseline laboratory characteristics. *Hypertension* 1991;17(Suppl 2):102-122.

Hertert S, Bailey G, Cottingham V, Carmody S, Egan D, Johnson P, Probstfield J. Community volunteers as recruitment staff in a clinical trial: the systolic hypertension in the elderly program (SHEP) experience [Published erratum appears in *Control Clin Trials* 1996;17:271]. *Control Clin Trials* 1996;17:23-32.

Kostis JB, Lawrence-Nelson J, Ranjan R, Wilson AC, Kostis WJ, Lacy CR, for the Systolic Hypertension in the Elderly (SHEP) Cooperative Research Group. Association of Increased Pulse Pressure With the Development of Heart Failure in SHEP. *Am J Hypertens* 2001;14:798-803.

Kostis JB, Davis BR, Cutler J, Grimm RH Jr., Berge KG, Cohen JD, Lacy CR, Perry HM Jr., Blaufox MD, Wassertheil-Smoller S, Black HR, Schron E, Berkson DM, Curb JD, Smith WM, McDonald R, Applegate WB. Prevention of heart failure by antihypertensive drug treatment in older persons with isolated systolic hypertension. SHEP Cooperative Research Group. *JAMA* 1997;278:212-216.

Kostis JB, Berge KG, Davis BR, Hawkins CM, Probstfield J. Effect of atenolol and reserpine on selected events in the systolic hypertension in the elderly program (SHEP). *Am J Hypertens* 1995;8:1147-1153.

Kostis JB, Lacy CR, Hall WD, Wilson AC, Borhani NO, Krieger SD, Cosgrove NM. The

effect of chlorthalidone on ventricular ectopic activity in patients with isolated systolic hypertension. The SHEP Study Group. *Am J Cardiol* 1994;74:464-467.

Kostis JB, Allen R, Berkson DM, Curb JD, Davey J, Grimm RH, Hall WD, Lacy CR, Lee M, Miller S, et al. Correlates of ventricular ectopic activity in isolated systolic hypertension. SHEP Cooperative Research Group. *Am Heart J* 1994;127:112-121.

Hall WD, Davis BR, Frost P, Hoffmeier M, O'Brien JE, Pace S, Page L, Schneider KA, Stamler J. Systolic Hypertension in the Elderly Program (SHEP). Part 7: Baseline laboratory characteristics. *Hypertension* 1991;17(Suppl 2):II102-II22.

Labarthe DR, Blafox MD, Smith WM, Lacy CR, Schnaper H, LaBaw F, Mascioli S, Davey J, Lakatos E. Systolic Hypertension in the Elderly Program (SHEP). Part 5: Baseline blood pressure and pulse rate measurements. *Hypertension* 1991;17(Suppl 2):II62-II76.

Moye LA, Davis BR, Hawkins CM, Probstfield JL. Conclusions and implications of the systolic hypertension in the elderly program. *Clin Exp Hypertens* 1993;15:911-924.

Perry HM, Davis BR, Price TR, Applegate WB, Fields WS, Guralnik JM, Kuller L, Pressel S, Stamler J, Probstfield JL. Effect of treating isolated systolic hypertension on the risk of developing various types and subtypes of stroke: the Systolic Hypertension in the Elderly Program (SHEP). *JAMA* 2000;284:465-471.

Perry HM, Berge KG, Blafox MD, Davis BR, Grimm RH, McDonald R, Pressel S, Schron E, Smith WM, Vogt TM. The systolic hypertension in the elderly program. In: Leonetti G, Cuspidi C, editors. *Hypertension in the elderly*. Dordrecht, The Netherlands: Kluwer Academic Press; 1994. p.123-137.

Petrovitch H, Vogt TM, Berge KG, for the SHEP Cooperative Research Group. Isolated systolic hypertension: lowering the risk of stroke in older patients. *Geriatrics* 1992;47:30-32, 35-38.

Probstfield JL, Applegate WB, Borhani NO, Curb JD, Cutler JA, Davis BR, Furberg CD, Hawkins CM, Lakatos E, Page LB, et al. The Systolic Hypertension in the Elderly Program (SHEP): an intervention trial on isolated systolic hypertension. SHEP Cooperative Research Group. *Clin Exp Hypertens [A]* 1989;11:973-989.

Savage PJ, Pressel SL, Curb JD, Schron EB, Applegate WB, Black HR, Cohen J, Davis BR, Frost P, Smith W, Gonzalez N, Guthrie GP, Oberman A, Rutan G, Probstfield JL, Stamler J. Influence of long-term, low-dose, diuretic-based, antihypertensive therapy on glucose, lipid, uric acid, and potassium levels in older men and women with isolated systolic hypertension: The Systolic Hypertension in the Elderly Program. SHEP Cooperative Research Group. *Arch Intern Med* 1998;158:741-751.

Schron EB, Wassertheil-Smoller S, Pressel S. Clinical trial participant satisfaction: survey of SHEP enrollees. SHEP Cooperative Research Group. *Systolic Hypertension in the*

Elderly Program. *J Am Geriatr Soc* 1997;45:934-938.

SHEP Cooperative Research Group. Implications of the systolic hypertension in the elderly program. *Hypertension* 1993;21:335-343.

SHEP Cooperative Research Group. Prevention of stroke by antihypertensive drug treatment in older persons with isolated systolic hypertension. Final results of the Systolic Hypertension in the Elderly Program (SHEP). *JAMA* 1991;265:3255-3264.

SHEP Cooperative Research Group. Rationale and design of a randomized clinical trial on prevention of stroke in isolated systolic hypertension. *J Clin Epidemiol* 1988;41:1197-1208.

Vogt TM, Schron E, Pressel S, Wasserthiel-Smoller S, Eddleman EE, Miller S, Stamler J. Systolic Hypertension in the Elderly Program (SHEP). Part 3: Sociodemographic characteristics. *Hypertension* 1991;17(Suppl 2):24-34.

Wassertheil-Smoller S, Applegate WB, Berge K, Chang CJ, Davis BR, Grimm R Jr., Kostis J, Pressel S, Schron E. Change in depression as a precursor of cardiovascular events. SHEP Cooperative Research Group (Systolic Hypertension in the elderly). *Arch Intern Med* 1996;156:553-561.

Weiler PG, Camel GH, Chiappini M, Greenlick MR, Hughes GH, Luhr JC, Moye LA, Mungas D, Perron M, Peters MH, et al. Systolic Hypertension of the Elderly Program (SHEP). Part 9: Behavioral characteristics. *Hypertension* 1991;17(Suppl 2):152-161.

Wittes J, Davis B, Berge K, Cohen JD, Grimm RH Jr., Hawkins CM, Kuller L. Systolic Hypertension of the Elderly Program (SHEP). Part 10: Analysis. *Hypertension* 1991;17(Suppl 2):162-167.

Young JH, Klag MJ, Muntner P, Whyte JL, Pahor M, Coresh J. Blood pressure and decline in kidney function: findings from the Systolic Hypertension in the Elderly Program (SHEP). *J Am Soc Nephrol* 2002;13:2776-2782.